

Order	Scientific name		Weybourne	Beeston Regis	West Runton	East Runton	Cromer	Overstrand	Sidestrand	Trimingham	Mundesley	Bacton	Walcott	Ostend (Arctic Fresh-water Bed)	Ostend	Happisburgh site 1	Happisburgh site 3	Happisburgh undifferentiated	Eccles	Sea Palling	Corton	Pakefield (Cromerian MIS 17)	Pakefield (unspecified horizon)	Kessingland	Covehithe, Easton Bavents
-------	-----------------	--	-----------	---------------	-------------	-------------	--------	------------	------------	------------	-----------	--------	---------	---------------------------------	--------	--------------------	--------------------	------------------------------	--------	-------------	--------	------------------------------	---------------------------------	-------------	---------------------------

Jonathan Stewart Collection (<http://www.northfolk.org.uk/js/index.html>)

Harrison, D., Bates, P. and Clayden, J., On the occurrence of *Galemys kormosi* (Schreuder, 1940) in the British Lower Pleistocene. *Acta Theriologica*, vol 33, 26: 369-378. 1988

Lister, A.M., The stratigraphical interpretation of large mammal remains from the Cromer Forest-bed Formation. In Turner, C. (Ed), *The early Middle Pleistocene in Europe*. Balkema, 1996.

McWilliams, B., *The Vertebrate Fossils of the Cromer Forest-bed*. Norwich Castle Museum and Art Gallery. Reprinted 1975

Stuart, A.J., Vertebrate faunas from the early Middle Pleistocene. In Turner, C. (Ed), *The early Middle Pleistocene in Europe*. Balkema, 1996